

Yang-Mills Theory at 60: Milestones, Landmarks and Interesting Questions*

Ling-Lie Chau

Department of Physics, University of California, Davis, CA 95616, USA
chau@physics.ucdavis.edu
<http://www.physics.ucdavis.edu/>

On the auspicious occasion of celebrating the 60th anniversary of the Yang-Mills theory, and Professor Yang's many other important contributions to physics and mathematics, I will highlight the impressive milestones and landmarks that have been established in the last 60 years, as well as some interesting questions that are worthy of answers from future researches. The paper is written (without equations) for the interest of non-scientists as well as of scientists.

Keywords: Yang-Mills theory at 60.

Contents

1	Overview	79
2	Yang-Mills Theory: Milestones and Interesting Questions	81
3	The (Anti-)Self-dual Yang-Mills Fields: Landmarks and Interesting Questions	83
4	Concluding Remarks	87
	Acknowledgments	88
	Attachment A	89
	Attachment B	94
	Attachment C	100
	Attachment D	101
	References	102

1. Overview^a

The 1954 Yang-Mills (YM) paper,¹ that set forth the YM theory with YM equations, made at least the following three major advances, if not revolutions, in theoretical and mathematical physics.

*Invited contribution to the Proceedings of the Conference on 60 Years of Yang-Mills Gauge Theories, IAS, NTU, Singapore, 25–28 May 2015. Also by invitation an earlier version of the paper (with slight differences) was published in the Dec. 30, 2016 issue of International Journal of Modern Physics A, *Int. J. Mod. Phys. A* **30**, 1530068 (2015).

^aThe author would recommend that readers first read Secs. 1 (Overview) and 4 (Concluding Remarks) before reading Secs. 2 and 3, which cover a vast amount of information in highlight fashion, supplemented with references and ample sources for references for those who would like to dive into details.

First, the YM equations generalized the Maxwell equations² from the Abelian gauge field equations, in which the Maxwell fields do not interact with themselves, to the more general non-Abelian gauge field equations, in which the YM fields do interact with themselves. It is akin to Einstein's general relativity (GR) equations³ generalizing Newton's equations for gravity,⁴ though in different ways. The Maxwell equations are exact by themselves and the YM equations are generalizations, leaping conceptually from the simpler special case of the Maxwell equations. In contrast, Newton's equations for gravity are weak-field approximations of Einstein's GR equations, so they still work for our daily lives and for the motions of the planets, except for minor but measurable GR improvements to agree with Nature, e.g., in the workings of the global positioning system (GPS) and in the precession of the perihelion of Mercury.

Second, the YM equations (including the anti-self-dual and the self-dual YM equations) are beautiful nonlinear partial differential equations not only for physics but also for mathematics, which have inspired the development of fundamental physics as well as beautiful mathematics, as has been the case with the GR equations.

Third, the YM theory manifested the local gauge covariance, which was first established in Maxwell's equations for electromagnetic interactions, as a principle for more general interactions (later found to be the weak and strong interactions, see below). In contrast, the GR theory manifests the local coordinate covariance as the principle for gravitational interactions.

After decades of further theoretical development and experimental work, amazingly, Nature is found to make use of these theoretical advances for interactions among elementary matter fields.^b The $SU(3)$ Yang–Mills fields are the mediating force fields for strong interactions. The $SU(2)$ Yang–Mills fields are the mediating force fields for weak interactions, unifying in an intricate way with the $U(1)$ Maxwell fields to become the $U(1) \times SU(2)$ Maxwell–Yang–Mills (MYM) mediating force fields for the unified electroweak interactions. Nature surprisingly chooses to use the simplest types of groups,^c $U(1) \times SU(2) \times SU(3)$, for electroweak and strong interactions, three of the four major interactions known in Nature.

For the last thirty years of his life, Einstein had searched for, unsuccessfully, the answer to the question⁵: Does Nature have a unified theory for all interactions? As the end of 2015 is approaching, sixty years after his passing in 1955, the answer to his question is still yet to be found. However, he might be pleased to know the progress mankind has made toward answering his question. At the time of his passing, when Yang–Mills theory was less than one year old, among the four

^bThe experimentally established elementary matter fields, as of 2015, are the fields of quarks, leptons, and the Higgs boson of mass $125 \text{ GeV}/c^2$.

^cA group is a set of mathematical elements, together with an operation called group-multiplication, under which the group has an identity and every group element has its inverse. Groups are the second simplest mathematical structure, next to sets, yet are so prominently used by Nature in formulating laws of interactions.

experimentally established major interactions: electromagnetic, weak, strong, and gravity, only for electromagnetism had a quantum field theory been established: quantum electrodynamics (QED), with the quantum Maxwell fields being the U(1) “glues”. Now a quantum field theory has been established for electroweak interactions, quantum flavor dynamics (QFD), which unifies electromagnetic and weak interactions in an intricate way — the quantum Maxwell fields remain to be the U(1) “glues” as in QED and the quantum Yang–Mills fields are the SU(2) “glues”. Also, a quantum field theory has been established for strong interactions, quantum chromodynamics (QCD), in which the quantum Yang–Mills fields are the SU(3) “glues”. So Einstein’s question can be asked in more specific terms: Will we find that QFD and QCD (the combination of the two is called the Standard Model of particle physics) are parts of a grand unified theory (GUT)? Will we find the ultimate quantum description of Einstein’s general relativity (QGR)? Will we find the ultimate unified theory, The Unified Theory (TUT) or The Theory of Everything (TOE), of GUT and QGR? To know how these questions are answered will make life richer.^d

2. Yang–Mills Theory: Milestones and Interesting Questions

The milestones to be highlighted in this section, except for Milestone 4, have all been theoretically developed and experimentally established, and set forth in modern graduate textbooks on quantum field theories. See the authoritative volumes with comprehensive and exhaustive references by S. Weinberg,⁶ and the celebration volume for the 50th anniversary of the Yang–Mills theory edited by ’t Hooft,⁷ for which Attachment A provides a copy of its Title page, Preface and Contents. Therefore, I will be very brief in highlighting the milestones in this section without giving references (except a few) to original papers which Refs. 6 and 7 comprehensively and exhaustively provide.

It is interesting to note that among the sixty-one Nobel Prizes in Physics⁸ from 1954 to 2014, twenty-six are related to the making of the Standard Model, while the 2015 one honors the experiment that gave evidence for the need of going beyond the Standard Model (see Milestone 5).

Milestone 1. Quantum flavor dynamics QFD has been theoretically developed and experimentally established for electroweak interactions,^e unifying electromagnetic and weak interactions. In QFD the quantum SU(2) Yang–Mills fields are the quantum mediating force fields for weak interactions, i.e., the SU(2) “weak glues”, among elementary particles with SU(2) “weak charges”, while the quantum U(1)

^dNature has been kind to humans in allowing their brains to understand the complexity of Nature bit by bit over time and then bring the pieces of understanding together into larger structures of meaning. How long will this process continue? Will the human brain ever discover the totality of Nature (finite or infinite)?

^eThe letter F in QFD is to indicate the fact that quarks and leptons exist in three different types called “flavors”.

Maxwell fields remain, as in QED, to be the quantum U(1) “electromagnetic glues” among particles with U(1) “electromagnetic charges”. In this intricate quantum unification the Maxwell field quanta, photons or light, are massless (as in QED) and can travel large distance in space void of matter with electromagnetic properties. Thus we can see each other, the moon, the sun, stars and galaxies. On the other hand, the SU(2) Yang–Mills quanta gain masses, through the Higgs mechanism (see Milestone 4. below), to become the massive W^+ , W^- and Z^0 particles that can travel only very short distances.

Interestingly, among the four major interactions, **only** weak interactions, mediated by the SU(2) YM quanta, W^+ , W^- and Z^0 , violate the discrete symmetries P (parity) and CP (parity and charge conjugation), as well as T (time reversal), while CPT symmetry stays true in all interactions.

Milestone 2. Quantum chromodynamics QCD has been theoretically developed and experimentally established for strong interactions.^f

In QCD the quantum SU(3) Yang–Mills fields are the quantum mediating forces for strong interactions, the SU(3) “color glues” of quanta called gluons. They interact with quarks according to their strong-interaction charge, the SU(3) “color charge”. (The leptons and the Higgs particle are SU(3) “colorless”, and therefore do not participate in strong interactions).

The QCD quanta with the U(1) and the SU(2) electroweak charges and “flavors” can exist freely. In contrast, the QCD quanta with non-zero SU(3) “color charges” cannot exist freely. Or, they are “color confined” or “color enslaved”. They have to combine to form SU(3) “colorless” particles to exist freely. For example, protons and neutrons are SU(3) “colorless” particles formed from “colorful” quarks and gluons. The SU(3) “colorless” particles formed from “colorful” gluons are called glueballs, which are still under theoretical study⁹ and experimental search.¹⁰

Also unusual is that the binding forces from the gluons decrease as distances decrease (corresponding to increasing energies) between the quarks — so QCD is an asymptotically free theory, quite opposite to that of QED.¹¹ These make QCD manageable to calculate in high energies and in high temperatures.¹² This has implications for cosmological studies on the early universe and for high energy experiments.¹³

Milestone 3. Local gauge covariance has been established to be the underlying principle of how Maxwell and Yang–Mills fields act as “glues” among the basic matter fields^{14,15} — the Maxwell fields as the “electromagnetic glues” and the Yang–Mills fields as the “weak glues” as well as the “strong (color) glues”. This principle has been used to develop new theories, e.g., the supersymmetric Yang–Mills theories, SYM,¹⁶ and even for supersymmetric gravity, SGR.¹⁷

^fThe letter C in QCD is to indicate the fact that the “SU(3) gluons” have different SU(3) quantum identities called SU(3) “colors”, conjured up by the color of photons, the “U(1) gluons”.

Milestone 4. A Higgs particle (possibly the first of many) of the Higgs mechanism (the ubiquitous “molasses”),¹⁸ which produces the masses of quarks and the masses of W^+ , W^- and Z^0 (the $SU(2)$ Yang–Mills quanta in weak interactions), was finally observed in 2012,^{19,20} almost fifty years after the original theoretical proposals. It was an admirable triumph of the collaborative spirit in scientific research!

Milestone 5. Toward the end of the twentieth century, experiments established that the mass differences between any two of the three-flavored neutrinos are not zero, in contradiction to the all-zero-mass neutrinos in QFD of the Standard Model. This fact forces consideration of going beyond the Standard Model.^{21–23}

Interesting questions

- As highlighted above, Nature has made versatile uses of the Yang–Mills fields, but not for uses on the macroscopic scales like the Maxwell fields and the Newton–Einstein fields.²⁴ Can a precise mathematical physics explanation for this be found?²⁵
- Is there a deeper guiding principle for the Higgs mechanism? How many Higgs particles there are? Will the mass matrices of quarks and of leptons (which are generated by the presence of the Higgs fields) be derived, so hopefully a deeper understanding of the origin of CP violations in weak interactions will also follow?
- What will be established to be the Beyond Standard Model? Will the supersymmetric extensions²⁶ become experimentally observed realities?
- Will a grand unification theory, GUT, be found in which the electroweak and the strong interactions are unified?
- Will GR be established theoretically and experimentally as a quantum theory?

For a succinct and insightful perspective on the last three questions, see Witten.²⁷

3. The (Anti-)Self-dual Yang–Mills Fields: Landmarks and Interesting Questions

The Yang–Mills fields in even dimensions of spacetime can be expressed as a sum of two terms: the self-dual term and the anti-self-dual term. Fields that are self-dual (SDYM) or anti-self-dual (ASDYM) are special because they automatically satisfy the Yang–Mills equations, and in addition, by definition, satisfy nonlinear differential equations, one-order lower than the Yang–Mills equations, which are called the (A)SDYM equations. (A)SDYM fields are simpler to study, while still remaining interesting and important for researches in physics and mathematics.

Landmark 1. The (A)SDYM fields have interesting solutions called instantons, which possess specific topological numbers (the Pontryagin or the second-Chern number) and minimize the Yang–Mills action.

In 1975, Belavin, Polyakov, Schwartz and Tyupkin constructed explicitly the first such solutions,²⁸ which stimulated much interest in mathematics as well as in physics.

In 1977 Ward showed²⁹ that (A)SDYM fields are naturally described by the twistor formulation that Penrose developed³⁰ to describe massless fields and the self-dual Einstein fields, and derived what are now called the Penrose–Ward transformations. Not long afterwards, this breakthrough by Ward led to the full description of the space of instantons in 1979 by Atiyah, Drinfeld, Hitchin and Mannin (now called the ADHM construction).³¹ See the comprehensive review lecture by Atiyah and references to original papers.³²

The instanton solutions prompted intensive research on their non-perturbative effects in QCD and QFD,³³ and led to experimentally testable predictions, for example the axions³⁴ and the related sphaleron phenomena.³⁵

Landmark 2. For (A)SDYM fields in four-dimensions Yang showed³⁶ that (A)SDYM fields can be written as second-order nonlinear partial differential equations, now called the Yang equations, in terms of group-value local fields.[§]

Later the Yang equations were found to have many characteristics of the classical integrable systems in lower dimensions: Backlund transformations,³⁷ non-local conservation laws,³⁸ Riemann–Hilbert transformation properties,³⁹ Painleve properties,⁴⁰ generalized Riccati equations,⁴¹ and, most importantly, the linear systems (the Lax systems),⁴² and Kac-Moody algebras.^{43–46} For an overview of the work and references during this period of discoveries, see “*Integrable Systems*”,⁴⁷ the lectures therein^{48,49} and Attachment B below, which gives the memorable front pages of the book. For a glimpse of the landscape of mathematical physics in the late 1980s, see Chau and Nahm.⁵⁰

Additionally, many integrable systems in lower dimensions, including the famed Nahm equations,⁵¹ can be directly reduced from the Yang equations. See the in-depth reviews^{52,53} and the extensive references therein. See the overview⁵⁴ of (A)SDYM as a classical integrable system in four dimensions and its relations to those in lower and higher than four dimensions. The review Ref. 49 includes discussions on extensions to supersymmetric theories, SYM and SGR, which will be discussed in Landmark 4.

Therefore, (A)SDYM equations have been shown to be classical integrable systems in **four** dimensions that interestingly also serve as conceptual pathways between those integrable systems in dimensions lower and higher than four dimensions, as well as in supersymmetric dimensions.

The next important step was to develop (A)SDYM equations into a quantum integrable field theory in four dimensions. In the 1990s, the Yang equations were put to quantization, using the action constructed by Nair and Schiff⁵⁵ and by Hou

[§]Group-valued local fields are special. Usually group-valued fields are non-local, defined on a loop, not at points as local fields are.

and Song,⁵⁶ in terms of the Lie-valued fields^{55,57} and in terms of the group-valued local fields.⁵⁸ In Ref. 58 Yamanaka and I found that the quantized group-valued local fields are bi-module fields which satisfy intricate exchange algebras with structure coefficients satisfying generalized Yang–Baxter equations. So here Yang–Mills met Yang–Baxter for the first time with the quantum group properties,^{59,60} originated from the Yang–Baxter equations in low dimensions, appearing in a four-dimensional quantum theory for the first time. Further, the algebraic relations satisfied by the Lie-valued fields derived in Refs. 55 and 57 can be derived from the exchange algebras of the group-value local fields. For further developments see Popov–Preitschopf⁶¹ and Popov,⁶² in 1996 and in 1999 respectively.

However, the quantum contents of the Yang equations have not been fully developed in detail, and the full quantum versions of its classical integrability properties are yet to be revealed.

Landmark 3. Using (A)SDYM fields as tools, Donaldson^{63,64} made major discoveries about the four-manifolds.^h

Here was what Atiyah said on the occasion of the awarding of the 1986 Fields Medal to Donaldson,⁶⁵ referring to Donaldson’s paper Ref. 63 as [1]:

“In 1982, when he was a second-year graduate student, Simon Donaldson proved a result [1] that stunned the mathematical world. Together with the important work of Michael Freedman (described by John Milnor), Donaldson’s result implied that there are “exotic” 4-spaces, i.e., 4-dimensional differentiable manifolds which are topologically but not differentiably equivalent to the standard Euclidean 4-space R^4 . What makes this result so surprising is that $n = 4$ is the only value for which such exotic n -spaces exist. These exotic 4-spaces have the remarkable property that (unlike R^4) they contain compact sets which cannot be contained inside any differentiably embedded 3-sphere!

To put this into historical perspective, let me remind you that in 1958 Milnor discovered exotic 7-spheres, and that in the 1960s the structure of differentiable manifolds of dimension > 5 was actively developed by Milnor, Smale (both Fields Medalists), and others, to give a very satisfactory theory. Dimension 2 (Riemann surfaces) was classical, so this left dimensions 3 and 4 to be explored. At the last Congress, in Warsaw, Thurston received a Fields Medal for his remarkable results on 3-manifolds, and now at this Congress we reach 4-manifolds. I should emphasize that the stories in dimensions 3, 4, and $n > 5$ are totally different, with the low-dimensional cases being much more subtle and intricate.

The surprise produced by Donaldson’s result was accentuated by the fact that his methods were completely new and were borrowed from theoretical physics, in the form

^h A manifold is a space that can be covered by local patches that are like the Euclidean space, or the Minkowski space.

of the Yang–Mills equations. These equations are essentially a nonlinear generalization of Maxwell’s equations for electro-magnetism, and they are the variational equations associated with a natural geometric functional. Differential geometers study connections and curvature in fibre bundles, and the Yang–Mills functional is just the L^2 -norm of the curvature. If the group of the fibre bundle is the circle, we get back the linear Maxwell theory, but for nonabelian Lie groups, we get a nonlinear theory. Donaldson uses only the simplest non-Abelian group, namely $SU(2)$, although in principle other groups can and will perhaps be used.

Physicists are interested in these equations over Minkowski space–time, where they are hyperbolic, and also over Euclidean 4-space, where they are elliptic. In the Euclidean case, solutions giving the absolute minimum (for given boundary conditions at ∞) are of special interest, and they are called instantons.

Several mathematicians (including myself) worked on instantons and felt very pleased that they were able to assist physics in this way. Donaldson, on the other hand, conceived the daring idea of reversing this process and of using instantons on a general 4-manifold as a new geometrical tool. In this he has been brilliantly successful: he has unearthed totally new phenomena and simultaneously demonstrated that the Yang–Mills equations are beautifully adapted to studying and exploring this new field.

Donaldson’s works continue to be highly honored.⁶⁶

Then in 1994 Witten⁶⁷ showed that the Seiberg–Witten equations⁶⁸ of $N = 2$ SYM also provide a powerful tool for analyzing four-manifolds, confirming those by Donaldson and discovering new properties, which generated much excitement among mathematicians.⁶⁹

Landmark 4. The landmarks above resulting from studying the (A)SDYM fields have inspired extensions into supersymmetric theories.

In 1978, Witten generalized the concept advanced by Ward in Landmark 1,⁷⁰ that (A)SDYM equations are the results of certain integrability conditions,²⁹ to the full extent of YM and SYM. Of particular interest is that the integrability along light-like lines precisely gives the $N = 3$ SYM equations.¹⁶ This naturally led to the discovery of many classical integrability properties of the theory and also to the construction of group-valued local fields and the derivation of their equations, giving the generalized Yang equations of SYM.⁷¹

In 1986 Lim and I showed that $N = 5, 6, 7, 8$ SGR, in the linearized approximation, had a similar interpretation.⁷²

In 1992, I derived the linear systems for high- N SGR,⁷³ which signal the classical integrability properties, and constructed group-valued local fields and derived the equations they satisfy, giving the generalized Yang equations of SGR.

Recently, quantum $N = 4$ SYM and its quantum integrability properties have been studied very actively,⁷⁴ giving a renaissance to studies on S-matrices.^{75,76} For the current status of such studies, see the paper by Brink in this *Proceedings*.⁷⁷

So the landmarks of this section have revealed that (A)SDYM fields in four dimensions are important to study, both for physics and for mathematics. They also serve importantly as conceptual pathways between integrable systems in lower and higher dimensions, as well as to those in supersymmetric spaces.¹

Interesting questions

- What will be the experimental observations of the effects related to instantons?
- What will be the quantum manifestations of the classical integrability properties of the Yang equations? Do quantum bi-module fields have physical reality?
- Will experiments find that Nature uses supersymmetric theories, SYM and SGR?

If yes, such a huge discovery will inevitably lead us to ask whether SYM or SGR theories have quantum integrability properties?

- What are the implications to physics of the four-manifolds discovered by Donaldson?

4. Concluding Remarks

In the 60 years since the Yang–Mills paper, we have seen the establishment of quantum flavor dynamics, QFD, unifying electromagnetic and weak interactions into electroweak interactions, with Maxwell fields acting as the “U(1) glues” and Yang–Mills fields acting as the “SU(2) glues”. The quanta of the U(1) Maxwell fields, photons, are massless. They can travel far and are our light (in its full spectrum). In contrast, the quanta of SU(2) Yang–Mills fields for weak interactions gain masses from the Higgs mechanism and become massive W^+ , W^- and Z^0 particles. They can travel only for very short distances.

We have also seen the development of quantum chromodynamics QCD for strong interactions, with the SU(3) Yang–Mills fields acting as the “SU(3) color glues” and its quanta, the massless gluons, interacting among themselves and mediating interactions among quarks that have “SU(3) color charges”. Unlike particles with U(1) electric charges and/or SU(2) weak-interaction charges, the “SU(3) colored” gluons and quarks cannot exist freely. To be free they must form “SU(3) colorless” particles: glueballs made of gluons, hadrons (protons, neutrons, etc.) made of quarks and gluons, or be at super high energies or super high temperatures.

Nature chooses these versatile ways to use the Yang–Mills fields; however, it does not allow us to experience the Yang–Mills fields in our daily lives as we do the Maxwell fields and the Newton–Einstein fields. Why this is so is an interesting mathematical physics question worthy of an answer.

After the celebration of 60 years of the Yang–Mills Theory in 2014–2015 comes the celebration of 100 years of Einstein’s theory of general relativity in 2015–2016.

¹In Attachment C, I propose to add the topic of (A)SDYM fields to the list of the other thirteen topics that Professor Yang considered important and to which he made major contributions.

We are now being treated to progress reports about the advances made in our understanding of it,^{78,79} in particular its quantum theoretical description and experimental attempts to observe its possible quantum phenomena. As of now, near the end of 2015, no definitive quantum theoretical description for general relativity has been established and no quantum phenomena of general relativity have yet been observed — no gravitational waves (happily this statement needs to be revised on February 11, 2016),⁸⁰ not to mention the more illusive gravitons, their quanta.

The ultimate question is whether Nature makes use of a unified theory for all interactions.

With all these interesting questions, mankind will be kept busy intellectually for generations to come. We look forward to seeing what new discoveries will be celebrated when Yang–Mills theory turns 70 in 2024 and Einstein’s general relativity turns 110 in 2025.⁸¹


Acknowledgments

I thank Prof. Phua, Kok Khoo, the Founding Director of the Institute of Advanced Studies (IAS), Nanyang Technological University (NTU), Singapore, for his gracious invitation to me to participate in the momentous event of celebrating the 60th anniversary of the Yang–Mills theory at NTU in 2015 (see poster in Attachment D), and to contribute a paper for the Proceedings, and for his further invitation to publish the paper also in the *International Journal of Modern Physics A*. I appreciate very much the excellent work performed by the editorial and production team of World Scientific Publishing Co Pte Ltd, of which Prof. Phua is the Chairman and the Editor-in-Chief.

Also, I would like to give my heartfelt thanks to many friends, especially Ms. Karen Andrews, Prof. Nicolai (Kolya) Reshetikhin, Ms. Gloria Rogers, Prof. William (Bill) Rogers, and Mr. Weiben Wang, for reading the drafts and making helpful comments which improved the presentation of the paper, and for their warm friendship and encouragement that made the task of writing this paper more enjoyable; and offer my deep gratitude to Kolya, Professor of Mathematics who has served two multi-year terms as the editor of *Communications in Mathematical Physics* (Springer), and Bill, Professor of English, for their expert critiques that sharpened the paper. Of course, any errors in the paper are mine alone.

Attachment A

The Title Page, Preface and Contents of *50 Years of Yang-Mills Theory*, ed. G. 't Hooft (World Scientific, 2004).



Attachment A (Continued)

PREFACE

At the time of the publication of this volume, fifty years have passed since the appearance of an article in *The Physical Review* by Chen Ning Yang and Robert L. Mills, entitled “Conservation of Isotopic Spin and Isotopic Gauge Invariance”. This book on the one hand serves as a tribute to that monumental piece of work, and on the other intends to show how its subject has evolved since that time, highlighting the landmarks that followed after the original paper emerged, and allowing its authors to indulge in new ideas and concepts. Gauge Theory has indeed grown into a pivotal concept in the Theory of Elementary Particles, and it is expected to play an equally essential role in even more basic theoretical constructions that are speculated upon today, with the aim of providing an all-embracing picture of the universal Laws of Physics.

Some of the chapters in this book are contributions that have appeared elsewhere; most of the contributions are original pieces of work. All are accompanied by brief comments by the Editor. Needless to state that this volume is far from complete. There are numerous well-known landmarks that we could not cover. Furthermore, like most developments in Science, progress not only comes from the relatively small set of papers by famous authors that enjoy enormous scores on citation indices, but it predominantly comes from the large crowds of scientists who confirm and reproduce the original research while adding inconspicuous but essential bits of understanding, not only by writing papers, but also by lecturing to students, by performing experiments and doing calculations. Without them, this book could not have been written.

Attachment A (Continued)

CONTENTS	
<i>Preface</i>	v
Introduction 1	
<i>Chapter 1</i> Gauge Invariance and Interactions; Remembering Robert Mills C. N. Yang	7
Quantizing Gauge Field Theories 13	
<i>Chapter 2</i> The Space of Gauge Fields: Its Structure and Geometry Bryce DeWitt	15
Ghosts for Physicists 33	
<i>Chapter 3</i> Perturbation Theory for Gauge-Invariant Fields V. N. Popov and L. D. Faddeev	39
Breaking the Symmetry 61	
<i>Chapter 4</i> Broken Symmetry and Yang–Mills Theory François Englert	65
Towards the Standard Model 97	
<i>Chapter 5</i> The Making of the Standard Model Steven Weinberg	99

Attachment A (Continued)


viii	50 Years of Yang–Mills Theory	
	Renormalization	119
<i>Chapter 6</i>		
Renormalization		121
Gerardus 't Hooft		
<i>Chapter 7</i>		
From Koszul Complexes to Gauge Fixing		137
R. Stora		
<i>Chapter 8</i>		
The Renormalization Content of Slavnov–Taylor Identities		168
Carlo Becchi		
	Anomalies	183
<i>Chapter 9</i>		
Anomalies to All Orders		187
Stephen L. Adler		
<i>Chapter 10</i>		
Fifty Years of Yang–Mills Theory and Our Moments of Triumph		229
Roman Jackiw		
	Asymptotic Freedom	253
<i>Chapter 11</i>		
Yang–Mills Theory In, Beyond, and Behind Observed Reality		255
Frank Wilczek		
	Magnetic Monopoles	269
<i>Chapter 12</i>		
To Be Or Not to Be?		271
F. Alexander Bais		
	Quark Confinement and Strings	309
<i>Chapter 13</i>		
Confinement and Liberation		311
A. M. Polyakov		

Attachment A (Continued)

<i>Contents</i>	<i>ix</i>
Fixing in Gauge Condition Non-Perturbatively 331	
<i>Chapter 14</i>	
Non-Perturbative Aspects of Gauge Fixing	333
Pierre van Baal	
The Lattice 355	
<i>Chapter 15</i>	
Yang–Mills Fields and the Lattice	357
Michael Creutz	
Fermions on the Lattice 375	
<i>Chapter 16</i>	
Chiral Symmetry on the Lattice	377
P. Hasenfratz	
Confrontation with Experiment 399	
<i>Chapter 17</i>	
Fifty Years of Yang–Mills Theories: A Phenomenological Point of View	401
Alvaro De Rújula	
Supersymmetry and Supergravity 431	
<i>Chapter 18</i>	
Supergravity as a Yang–Mills Theory	433
Peter van Nieuwenhuizen	
Physics of the 21st Century 457	
<i>Chapter 19</i>	
Gauge/String Duality for Weak Coupling	459
Edward Witten	

Attachment B

The Title Page, Forewords by S. S. Chern and C. N. Yang, and Contents of *Integrable Systems*, Nankai University, Tianjin, China; ed. X. C. Song, (World Scientific, 1987).



Attachment B (Continued)

數學物理

永結一體

陳省身



一九八八年九月

Mathematics and Physics
One Body Forever

S.S.Chern

Attachment B (Continued)

数学物理是很古老的学科.近年来发展
尤其迅速.南开数学所理论物理研究室
将出版一系列关于此学科的书籍,这是
很有意义的事.

楊振寧

Mathematical physics is a subject of
study which dated far back to ancient times,
and has found speedy development in recent
years.

It is most significant that the Theore-
tical Physics Division of Nankai Institute
of Mathematics will publish a series of books
on the subject.

C.N.Yang

Attachment B (Continued)

CONTENTS	
INSCRIPTIONS	IV
PREFACE	IX
SPEECHES	
On the Relation between Mathematics and Physics <i>L.D.Faddeev</i>	3
Journey through Statistical Mechanics <i>C.N.Yang</i>	11
LECTURES	
Lectures on Quantum Inverse Scattering Method <i>L.D.Faddeev</i>	23
Lecture I	24
Lecture II	34
Lecture III	48
Lecture IV	59
Geometrical Integrability and Equation of Motion in Physics: A Unifying View <i>L.L.Chau</i>	71
CONTRIBUTIONS	
Generalization of Sturm-Liouville Theory <i>C.N.Yang</i>	154
Generalized Levinson's Theorem <i>Z.Q.Ma</i>	155

Attachment B (Continued)

xii	
On the Darboux Form of Bäcklund Transformations	162
<i>C.H.Gu</i>	
A Trace Identity and Its Applications in Hamiltonian Theory of Integrable Systems	169
<i>G.Z.Tu</i>	
On the Hierarchies of Integrable Systems in 2+1 Dimensions	177
<i>Y.Cheng</i>	
The Relationships between Inverse Scattering Techniques and Conformal Symmetry Transformations in Certain Nonlinear Systems	183
<i>B.Y.Hou and W.Li</i>	
Hamiltonian Structures of the Two Dimensional Classical Integrable System, Loop Algebras and Basic Poisson Brackets	192
<i>M.L.Ge and J.F.Lü</i>	
Nonlinear Realization of Hidden Symmetry $U(1)$ and the Kac-Moody Algebra of Chiral Field in Curved Space	206
<i>X.C.Song</i>	
The Bosonization of (1+1) Dimensional Fermion Fields by Path Integration Method	220
<i>G.J.Ni, R.T.Wang and S.Q.Chen</i>	
Bethe Ansatz Equations for Multicomponent Nonlinear Schrödinger Model	226
<i>F.C.Pu and B.H.Zhao</i>	
LIST OF PARTICIPANTS	245

Attachment B (Continued)

77

WORKSHOP ON GEOMETRICAL INTEGRABILITY
AND EQUATIONS OF MOTION IN PHYSICS

	<u>Aug. 18</u>	<u>Aug. 19</u>	<u>Aug. 20</u>	<u>Aug. 21</u>	<u>Aug. 22</u>
8:30-9:30	Chau	Chau	Chau	Chau	Chau
10:00-11:00	"	"	"	"	"
Lunch					
2:30-3:30	Song	Yan	Ni	Contribu-	Panel
4:00-5:00	Hou	Ge		tions	Discussion
				"	

Topics:

- L.-L. Chau (乔玲丽), University of California, Davis
- I. Generic properties of geometrically integrable systems:
Flow Chart, and two dimensional systems;
 - II. Self-dual Yang-Mills Equations;
 - III. Supersymmetric Yang-Mills systems;
 - IV. Supergravity;
 - V. Summary and outlook.
- M.-L. Ge (葛墨林), Lanchou University and Nankai University, Tianjing
Integrability of Belinski-Zakharov Gravity and Solitons in
Gravitational Wave.
- B.-Y. Hou (侯伯宇), Xibei University, Xian
Virasoro and Kac-Moody of Ernst equation.
- G.-J. Ni (倪光炯), Fudan University, Shanghai
A general method of calculation of anomalies.
- X.-C. Song (宋行长), Beijing University, Beijing
Integrability properties of Ernst equation.
- M.-L. Yan (闫沐霖), Science and Technology University, Hofei
Potts Model, Yang-Baxter Equation's Solutions; Surfaces with Genus > 1 .

Attachment C

C. N. Yang's Commandments in Physics

When celebrating Professor Yang's 90th birthday, 2012, Tsinghua University presented him a black rock cube with his thirteen important contributions to physics listed and carved on the vertical faces, grouped into four major categories in physics (one on each vertical face of the cube), which I list in the table below (the first thirteen in black letters).


(The two photos of the cube below are copies of [Ph 72] and [Ph 73] in C. N. Yang, *Selected Papers II with Commentaries* (World Scientific, 2013). In the table below, I adopt the same numbering system and English translation as given by Yu Shi, "Beauty and physics: 13 important contributions of Chen Ning Yang," *Int. J. Mod. Phys. A* **29**, 1475001 (2014).)

Face-A: Statistical Mechanics: A.1. 1952 Phase Transition A.2. 1957 Bosons A.3. 1967 Yang-Baxter Equation A.4. 1969 Finite Temperature	Face-B: Condensed Matter Physics: B.1. 1961 Flux Quantization B.2. 1962 ODLRO
Face-C: Particle Physics: C.1. 1956 Parity Nonconservation C.2. 1957 T, C and P C.3. 1960 Neutrino Experiment C.4. 1964 CP Nonconservation	Face-D: Field Theory: D.1. 1954 Gauge Theory D.2. 1974 Integral Formalism D.3. 1975 Fiber Bundle D.4. 1977 (Anti-)Self-dual Gauge Fields

Here I would propose to call the listed important contributions


C. N. Yang's Commandments in Physics

and to add "1977 (Anti-) Self-dual Gauge Fields" to the list on Face-D (as D.4., shown in red) so to become the Fourteenth Commandment.



Attachment D

The poster of the Conference on 60 Years of Yang-Mills Gauge Theories, IAS, NTU, Singapore, 25–28 May 2015.



**NANYANG
TECHNOLOGICAL
UNIVERSITY**

Institute of Advanced Studies

Sponsored by

金基氏李
LEE FOUNDATION

Conference on 60 Years of Yang-Mills Gauge Field Theories

CN Yang's Contributions to Physics

25 to 28 May 2015 • Nanyang Technological University, Singapore

CO-CHAIRS

Lars BRINK Chalmers Univ. of Technology

Kok Khoo PHUA IAS, Nanyang Technological Univ.

ADVISORY BOARD

Sir Michael ATIYAH The Univ. of Edinburgh
Fields Medalist 1966

David GROSS Univ. of California, Santa Barbara
Nobel Laureate in Physics 2004

Ludwig FADDEEV Steklov Institute of Mathematics, RAS

Peter GODDARD IAS, Princeton

Roman JACKIW MIT

INTERNATIONAL ORGANISING COMMITTEE

Lay Nam CHANG Virginia Tech

Ngee Pong CHANG City College of New York & NTU, Singapore

Yongmin CHO Konkuk Univ.

Da Hsuan FENG Univ. of Macau

Kazuo FUJIKAWA RIKEN

Molin GE Nankai Univ.

Hua Zhong LI Sun Yat-sen Univ.

Zhong Qi MA Institute of High Energy Physics, CAS

Bruce MCKELLAR CoEPP & Univ. of Melbourne

Antti NIEMI Uppsala Univ.

George STERMAN Stony Brook Univ.

Edward WITTEN IAS, Princeton

Tai Tsun WU Harvard Univ.

Kenneth YOUNG The Chinese Univ. of Hong Kong

Bang Fen ZHU IAS, Tsinghua Univ.

SPEAKERS

David GROSS Univ. of California, Santa Barbara
Nobel Laureate in Physics 2004

Lars BRINK Chalmers Univ. of Technology

Hong Mo CHAN Rutherford Appleton Laboratory, UK

Lay Nam CHANG Virginia Tech

Ngee Pong CHANG City College of New York & NTU, Singapore

Alexander CHAO Stanford National Accelerator Laboratory

Ling Lie CHAU Univ. of California, Davis

Paul CHU Univ. of Houston

Robert CREASE Stony Brook Univ.

Michael CREUTZ Brookhaven National Laboratory

Totru EGUCHI Rikkyo Univ.

Ludwig FADDEEV Steklov Institute of Mathematics, RAS

Michael FISHER Univ. of Maryland

Kazuo FUJIKAWA RIKEN

Xi Wen GUAN Wuhan Institute of Physics and Mathematics, CAS

Sen HU Univ. of Science and Technology of China

Henrik JOHANSSON Uppsala Univ.

Louis KAUFFMAN Univ. of Illinois at Chicago

Ren Bao LIU Chinese Univ. of Hong Kong

Chen Ning YANG Tsinghua Univ.
Nobel Laureate in Physics 1957

Zhong Qi MA Institute of High Energy Physics, CAS

Bruce MCKELLAR CoEPP & Univ. of Melbourne

Antti NIEMI Uppsala Univ.

Benni REZNIK Tel Aviv Univ.

George SAVVIDY Institute of Nuclear and Particle Physics, Demokritos

Samson SHATASHVILI Trinity College Dublin

George STERMAN Stony Brook Univ.

Sheung Tsun TSOU Univ. of Oxford

Sze Hoi Henry TYE IAS, The Hong Kong Univ. of Science and Technology

Jie WEI Michigan State Univ.

Sau Lan WU Univ. of Wisconsin-Madison

Tai Tsun WU Harvard Univ.

Yong Shi WU Univ. of Utah & Fudan Univ.

Li Hua YU Brookhaven National Laboratory

Anthony ZEE Univ. of California, Santa Barbara


Hui ZHAI IAS, Tsinghua Univ.

Bang Fen ZHU IAS, Tsinghua Univ.

& others

During the last six decades, Yang-Mills theory has increasingly become the cornerstone of theoretical physics. It is seemingly the only fully consistent relativistic quantum many-body theory in four spacetime dimensions. As such it is the underlying theoretical framework for the Standard Model of Particle Physics, which has been shown to be the correct theory at the energies we now can measure. It has been investigated also from many other perspectives, and many new and unexpected features have been uncovered from this theory. In recent decades, apart from high energy physics, the theory has been actively applied in other branches of physics, such as statistical physics, condensed matter physics, nonlinear systems, etc. This makes the theory an indispensable topic for all who are involved in physics.

www.ntu.edu.sg/ias



CN Yang and Robert Mills

Supported by:



School of Physical and Mathematical Sciences



South East Asia
Theoretical Physics
Association

References

1. C. N. Yang and R. Mills, Isotopic spin conservation and a generalized gauge invariance, *Phys. Rev.* **95**, 631 (1954).
2. J. M. Maxwell, A dynamical theory of the electromagnetic field, *Philos. Trans. R. Soc. London* **155**, 459 (1865), doi:10.1098/rstl.1865.000 (This article followed a December 8, 1864 presentation by Maxwell to the Royal Society.)
3. A. Einstein, Feldgleichungen der gravitation (English translation: The field equations of gravitation) *Preussische Akademie der Wissenschaften, Sitzungsberichte*, 844–847, part 2 (1915); according to A. Pais of reference below (Chap. 14, Ref. E1; Chap. 15, Ref. E15), this is the defining paper of general relativity (GR) after several earlier publications on GR in 1915, which was then followed by many more in the following years, see https://en.wikipedia.org/wiki/List_of_scientific_publications_by_Albert_Einstein#Journal_articles.
4. I. Newton, *Philosophiæ Naturalis Principia Mathematica*, three books in Latin, first published 5 July 1687, often referred to as *The Principia*; English translation: *The Mathematical Principles of Natural Philosophy* (1729).
5. A. Pais, *Subtle Is the Lord: The Science and the Life of Albert Einstein* (Oxford University Press, 1982).
6. S. Weinberg, *The Quantum Theory of Fields Vol. I: Foundations* (Cambridge University Press, 1995); S. Weinberg, *The Quantum Theory of Fields Vol. II: Modern Applications* (Cambridge University Press, 1996); S. Weinberg, *The Quantum Theory of Fields Vol. III: Supersymmetry* (Cambridge University Press, 2000).
7. G. 't Hooft (ed.), *50 Years of Yang–Mills Theory* (World Scientific, Singapore, 2005).
8. For list of all Nobel Prizes in Physics, see <http://www.nobelprize.org/nobel-prizes/physics/laureates/>.
9. M. Creutz, The lattice and quantized Yang–Mills theory, in this *Proceedings*.
10. W. Ochs, The status of glueballs, *J. Phys. G: Nucl. Part. Phys.* **40**, 67 (2013), arXiv:1301.5183v3; F. Close and P. R. Page, Glueballs, *Sci. Amer.* **279**, 80 (1998).
11. D. J. Gross, Quantum chromodynamics: The perfect Yang–Mills gauge field theories, in this *Proceedings*; F. Wilczek, Yang–Mills theory in, beyond, and behind observed reality, in Ref. 7.
12. G. Sterman, Yang–Mills theories at high energy accelerators, in this *Proceedings*; R. Hofmann, *The Thermodynamics of Quantum Yang–Mills Theory: Theory and Applications* (World Scientific, 2012); *SU(2)* Yang–Mills thermodynamics, talk at the Conference.
13. CMS Collab. (V. Khachatryan *et al.*), Evidence for collective multiparticle correlations in p-Pb collisions, *Phys. Rev. Lett.* **115**, 012301 (2015); and news <http://earthsky.org/human-world/lhc-creates-liquid-from-big-bang>, LHC creates liquid from Big Bang, earthsky.org/human-world, Sep 15, 2015.
14. C. N. Yang, The conceptual origins of Maxwell's equations and gauge theory, *Phys. Today* **67**, 45 (2014).

15. C. N. Yang, Gauge invariance and interactions, in Ref. 7. It was a copy of his notes hand-written in 1947 when he was a graduate at University of Chicago — impressive that a graduate student would think about such a big question! A typed transcript of the notes is given in Attachment-1 of L. L. Chau’s “Almost everything Yang touched eventually turned into gold”, in *Proceedings of the Conference in Honor of C. N. Yang’s 85th Birthday*, eds. C. H. Oh, K. K. Phua, N. P. Chang and L. C. Kwek (World Scientific, 2008), <http://chau.physics.ucdavis.edu/Yang85-byChau-fromWS081104f.pdf>, Also Attachment-2 of this Chau’s paper gives an anecdote about Wigner’s attempt to understand gauge potential.
16. L. Brink, J. H. Schwarz and J. Scherk, Supersymmetric Yang–Mills theories, *Nucl. Phys. B* **121**, 77 (1977); G. Gliozzi, J. Scherk and D. Olive, Supersymmetry, supergravity theories and the dual spinor model, *Nucl. Phys. B* **122**, 253 (1977); and references therein for earlier papers.
17. P. van Nieuwenhuizen, Supergravity as a Yang–Mills theory, in Ref. 7; D. Z. Freedman and A. Van Proeyen, *Supergravity* (Cambridge University Press, 2012).
18. P. W. Higgs, Spontaneous symmetry breakdown without massless bosons, *Phys. Rev.* **145**, 1156 (1966); F. Englert, R. Brout and M. F. Thiry, Vector mesons in presence of broken symmetry, *Nuovo Cimento A* **43**(2), 244 (1966); For an overview and more references, see http://www.nobelprize.org/nobel_prizes/physics/laureates/2013/advanced-physicsprize2013.pdf.
19. ATLAS Collab., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012), [arXiv:1207.7214](https://arxiv.org/abs/1207.7214).
20. CMS Collab., Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
21. Particle Data Group (S. Raby), Grand unified theories (2011), <http://pdg.lbl.gov/2015/reviews/rpp2015-rev-guts.pdf>; See also Wikipidia, *Physics Beyond the Standard Model*, https://en.wikipedia.org/wiki/Physics_beyond_the_Standard_Model.
22. The leaders of the experiments are honored by the 2015 Nobel Prize in Physics, http://www.nobelprize.org/nobel_prizes/physics/laureates/2015/ which gives an overview and references about neutrino oscillation at https://www.nobelprize.org/nobel_prizes/physics/laureates/2015/advanced-physicsprize2015.pdf; They are also honored by the Breakthrough Prize, <https://breakthroughprize.org/>, in the category of Fundamental Physics 2016, https://en.wikipedia.org/wiki/Fundamental_Physics_Prize.
23. Both Advanced Information for the 2015 and for the 2008 Nobel Prizes in Physics, https://www.nobelprize.org/nobel_prizes/physics/laureates/2015/advanced-physicsprize2015.pdf and https://www.nobelprize.org/nobel_prizes/physics/laureates/2008/advanced-physicsprize2008.pdf have used the parameterization for the flavor mixing matrix originally given by L. L. Chau and W. Y. Keung, Comments on the parameterization of the Kobayashi-Maskawa matrix, *Phys. Rev. Lett.* **53**, 1802 (1984), <http://chau.physics.ucdavis.edu/Chau-Keung-paraKM-PRL53-p1802-1984.pdf> which has also been adopted by the Particle Data Group as the “standard choice”, <http://pdg.lbl.gov/2015/reviews/rpp2015-rev-ckm-matrix.pdf>, Ref. 3, after an interesting evolution of

choices. More importantly, the parameters of the parameterization which is in product form and in terms of exact variables (not approximate) have physical meaning in experiments (such as those honored by the 2015 Nobel Prize in Physics). They are precisely used by Nature to characterize flavor oscillations among quarks and among neutrinos, as well as to characterize CP violations. While CP violation in weak interactions of quark flavor changing has been established (after a long interesting history of theoretical and experimental work), CP violation or no CP violation in neutrino flavor changing is yet to be settled by experiments. It will surely be a milestone in the history of physics when that is settled, and then the parameterization will once again be highlighted. [Right now the data are consistent with three flavors of quarks and three flavors of leptons. However, in the event we are surprised in the future and find more than three flavors, the Chau-Keung type of parameterizations (in product form and in terms of exact variables) for any N flavors have already been derived and given in L. L. Chau, Phase direct CP violations and general mixing matrices, *Phys. Lett. B* **651**, 293–297 (2007), <http://chau.physics.ucdavis.edu/Chau-Direct-CP-PLb651-p293.pdf>].

Another close encounter with Nobel-Prize-honored discoveries was through the paper by R. F. Peierls, T. L. Trueman and L. L. (Chau) Wang, Estimates of production cross sections and distributions for W bosons and hadrons jets in high energy pp and $p\bar{p}$ collisions, *Phys. Rev. D* **16**, 1397 (1977). The results in this paper for the quantities given in the title agreed with those of the 1983 experiments in which the observation of W^+ , W^- and Z^0 earned the 1984 Nobel Prize in Physics for C. Rubbia and S. van der Meer, <http://www.nobelprize.org/nobel-prizes/physics/laureates/1984/>. After the suspension of several months between the observation of the W 's and the observation of Z^0 , amazingly (or boringly, depending on one's point of view) the masses of W^+ , W^- and Z^0 were observed at the predicted values based upon the earlier 1979 Nobel-Prize-honored theoretical works by S. L. Glashow, by A. Salam and by S. Weinberg, <http://www.nobelprize.org/nobel-prizes/physics/laureates/1979/>. This success story concurs once more with Einstein's famous statement, "*Subtle is The Lord, but malicious He is not!*" (The English translation from German is that of A. Pais, Ref. 5.)

The reason that Peierls, Trueman and (Chau) Wang did the calculations reported in that paper was because Brookhaven National Laboratory (BNL or Brookhaven) was building the then highest energy pp (proton-proton) collider, ISABELLE, in search of W^+ , W^- and Z^0 . Unfortunately Brookhaven failed to produce magnets with the required properties. This resulted in the cancellation of ISABELLE, after already spending \$200M, <https://en.wikipedia.org/wiki/ISABELLE> (so goes the joke, "IS-A-BELLE became WAS-A-BELLE"). As a consequence, leading experimental physicists left BNL and went to CERN, Geneva, Switzerland to build the proton-anti-proton $p\bar{p}$ collider and then made the observation of W^+ , W^- and Z^0 . (As the authors of the paper we were told by the experimentalists that the results of substantial $p\bar{p}$ production cross sections for W^+ , W^- and Z^0 in our paper gave them confidence.) The momentous event also planted the seed that eventually led to the building of the now world-famed Large Hadron Collider LHC at Geneva, where the discovery of the Higgs particle was made in 2012, Refs. 18, 19, 20. Brookhaven had lost such a golden chance. One can only imagine what it would be like for Brookhaven and for US

- high energy physics had Brookhaven been successful in building ISABELLE.
24. This is related to the problem of “Yang–Mills and mass gap”, which is the first of the seven Millennium Problems posted by the Clay Mathematics Institute <http://www.claymath.org/millennium-problems>, with official problem description at <http://www.claymath.org/sites/default/files/yangmills.pdf>, and the status as of 2004 at <http://www.claymath.org/sites/default/files/ym2.pdf>.
 25. E. Witten, The problem of gauge theory, <http://arxiv.org/pdf/0812.4512v3.pdf>, where he gave the following concluding statements: “*The mass gap is the reason, if you will, that we do not see classical nonlinear Yang–Mills waves. They are a good approximation only under inaccessible conditions. I have spent most of my career wishing that we had a really good way to quantitatively understand the mass gap in four-dimensional gauge theory. I hope that this problem will be solved one day.*” See also his, *What We Can Hope to Prove About 3d Yang–Mills Theory*, http://media.scgp.stonybrook.edu/presentations/20120117_3_Witten.pdf Simons Center, January 17, 2012.
 26. S. Weinberg, *Supersymmetry, Vol. III*, of Ref. 6; J. Terning, *Modern Supersymmetry: Dynamics and Duality* (Clarendon Press, Oxford, 2006).
 27. E. Witten, Quest for unification, <http://arxiv.org/pdf/hep-ph/0207124.pdf>, based on his Heinrich Hertz lecture at SUSY 2002 at DESY, June, 2002.
 28. A. A. Belavin, A. M. Polyakov, A. S. Schwartz and Yu. S. Tyupkin, Pseudoparticle solutions of the Yang–Mills equations, *Phys. Lett. B* **59**, 85 (1975).
 29. R. S. Ward, On self-dual gauge fields, *Phys. Lett. A* **61**, 81 (1977).
 30. R. Penrose, Zero rest-mass fields including gravitation: Asymptotic behaviour, *Proc. R. Soc. A* **284**, 159 (1965); The twistor programme, *Rep. Math. Phys.* **12**, 65 (1977).
 31. M. F. Atiyah, V. G. Drinfeld, N. J. Hitchin and Yu. I. Manin, Construction of instantons, *Phys. Lett. A* **65**, 185 (1978).
 32. M. F. Atiyah, *Geometry of Yang–Mills Fields* (Publications of the Scuola Normale Superiore, Pisa, 1979).
 33. See the discussions and references on instantons in Chapter 23, Vol. II. of Ref. 6.
 34. M. Kuster, G. Raffelt and B. Beltrán (eds.), *Axions: Theory, Cosmology, and Experimental Searches*, Lecture Notes in Physics, Vol. 741 (Springer-Verlag, Berlin, Heidelberg, 2008).
 35. H. Tye, Sphaleron physics in Yang–Mills theory, talk at the Conference.
 36. C. N. Yang, Conditions of self-duality for SU(2) gauge fields on Euclidean four-dimensional space, *Phys. Rev. Lett.* **38**, 1377 (1977).
 37. M. K. Prasad, A. Sinha and L. L. Chau Wang, Parametric Backlund transformation for self-dual SU(N) Yang–Mills fields, *Phys. Rev. Lett.* **43**, 750 (1979).
 38. M. K. Prasad, A. Sinha and L. L. Chau Wang, Non-local continuity equations for self-dual SU(N) Yang–Mills fields, *Phys. Lett. B* **87**, 237 (1979).
 39. K. Ueno and Y. Nakamura, Transformation theory for anti-(self)-dual equations and the Riemann–Hilbert problem, *Phys. Lett. B* **109**, 273 (1982).
 40. R. S. Ward, The Painleve properties for the self-dual gauge-field equations, *Phys. Lett. A* **102**, 279 (1984).

41. L. L. Chau and H. C. Yen, Generalized Riccati equations for self-dual Yang–Mills fields, *J. Math. Phys.* **28**, 1167 (1987).
42. L. L. Chau, M. K. Prasad and A. Sinha, Some aspects of the linear systems for self-dual Yang–Mills fields, *Phys. Rev. D* **24**, 1574 (1981).
43. L. L. Chau, M. L. Ge and Y. S. Wu, Kac-Moody algebra in the self-dual Yang–Mills equations, *Phys. Rev. D* **25**, 1086 (1982); L. L. Chau, M. L. Ge, A. Sinha and Y. S. Wu, Hidden-symmetry algebra for the self-dual Yang–Mills equation, *Phys. Letts. B* **121**, 391 (1983).
44. L. Dolan, Kac-Moody symmetry group of real self-dual Yang–Mills, *Phys. Lett. B* **113**, 378 (1982).
45. L. L. Chau and Y. S. Wu, More about hidden-symmetry algebra for the self-dual Yang–Mills system, *Phys. Rev. D* **26**, 3581 (1982), see the discussion of comparing results with Ref. 44.
46. L. Dolan, Kac-Moody algebras and exact solvability in hadronic physics, *Phys. Rep.* **109**, 1 (1984).
47. X. C. Song (ed.), *Integrable Systems* (World Scientific, 1987), (Nankai University, Tianjin, China).
48. L. D. Faddeev, Lectures on quantum inverse scattering method, in Ref. 47.
49. L. L. Chau, A unifying description of equations of motion in physics from geometrical integrability, in Ref. 47.
50. L. L. Chau and W. Nahm (eds.), *Differential Geometric Methods in Physics: Physics and Geometry, Proceedings of the NATO Advanced Research Workshop and the 18th International Conference*, University of California, Davis, 1988 (Plenum Press, 1990); and see its Title Pages and Contents at <http://chau.physics.ucdavis.edu/Text/pdfs/NATO-Intl-Conf-Diff-Geo-Th-Phys-UCD-1988-Jul2-8.pdf>.
51. W. Nahm, The construction of all self-dual multimonopoles by the ADHM method, in *Monopoles in Quantum Field Theory* (World Scientific, Singapore, 1982), pp. 87–94, See also F. Alexander Bais, Magnetic monopoles in non-Abelian gauge theories, in Ref. 7.
52. L. J. Mason and N. M. J. Woodhouse, *Integrability, Self-Duality, and Twistor Theory*, LMS Monograph, New Series, Vol. 15 (Oxford University Press, Oxford, 1996).
53. M. J. Ablowitz, S. Chakravarty and R. G. Halburd, Integrable systems and reductions of the self-dual Yang–Mills equations, *J. Math. Phys.* **44**, 3147 (2003).
54. R. S. Ward, Integrable and solvable systems, and relations among them, *Phil. Trans. R. Soc. London A* **315**, 451 (1985).
55. V. E. Nair and J. Schiff, A Kähler-Chern-Simons theory and quantization of instanton moduli, *Phys. Lett. B* **246**, 423 (1990); Kahler Chern–Simons theory and symmetries of anti-self-dual gauge fields, *Nucl. Phys. B* **371**, 329 (1992).
56. B. Y. Hou and X. C. Song, Stony Brook ITP- SB-83-66, (1983), unpublished; Lagrangian form of the self-dual equations for $SU(N)$ gauge fields on four-dimension Euclidean space, *Commun. Theor. Phys.* **29**, 443 (1998).

57. A. Losev, G. Moore, N. Nekrasov and S. Shatashvili, Four-dimensional avatars of two-dimensional RCFT, *Nucl. Phys. B (Proc. Suppl.)* **46**, 130 (1996).
58. L. L. Chau and I. Yamanaka, Canonical formulation of the self-dual Yang–Mills system: Algebras and hierarchy, *Phys. Rev. Lett.* **68**, 1807 (1993); L. L. Chau and I. Yamanaka, Quantization of the self-dual Yang–Mills system: Exchange algebras, and local quantum group in four-dimensional quantum field theories, *Phys. Rev. Lett.* **70**, 1916 (1993).
59. V. G. Drinfeld, Hopf algebras and the quantum Yang–Baxter equation, *Dokl. Akad. Nauk, SSSR* **283**, 1060 (1985); Quantum groups, in *Proc. Int. Congr. Math. (Berkeley)*, Vol. I (Academic Press, New York, 1986), pp. 798–820; M. Jimbo, A q -analog of $U(\mathfrak{gl}(N + 1))$ Hecke algebras and the Yang–Baxter equation, *Lett. Math. Phys.* **11**, 247 (1986).
60. N. Yu. Reshetikin, L. A. Takhtadzhyan and L. D. Faddeev, Quantization of Lie groups and Lie algebras, *Leningrad Math. J.* **1**(1) (1990); M. Jimbo (ed.), *Yang–Baxter Equation in Integrable Systems*, Advanced Series in Mathematical Physics, Vol. 10 (World Scientific, 1990).
61. A. D. Popov and C. R. Preitschopf, Extended conformal symmetries of the self-dual Yang–Mills equations, *Phys. Lett. B* **374**, 71 (1996).
62. A. D. Popov, Holomorphic Chern–Simons–Witten theory: From 2D to 4D conformal field theories, *Nucl. Phys. B* **550**, 585 (1999).
63. S. K. Donaldson, Self-dual connections and the topology of smooth 4-manifolds, *Bull. Amer. Math. Soc.* **8**, 81 (1983).
64. S. K. Donaldson’s web page, <http://www.imperial.ac.uk/~skdona/>, gives web links to all his publications; complete at <http://www.ams.org/mathscinet/>, and since 2004 at <http://arxiv.org/>.
65. M. Atiyah, On the work of Simon Donaldson, in *Proceedings of the International Congress of Mathematicians*, Berkeley, California, USA, 1986.
66. See the profile of S. K. Donaldson at https://en.wikipedia.org/wiki/Simon_Donaldson, The citation of his work for his 1986 Fields Medal at https://en.wikipedia.org/wiki/Fields_Medal, The citation of his work for the 2009 Shaw Prizes in Mathematical Sciences at <http://www.shawprize.org/en/shaw.php?tmp=3&twoid=12&threeid=41&fourid=20>, and see him as one of the seven recipients of the inaugural Breakthrough Prizes in Mathematics, 2014, https://en.wikipedia.org/wiki/Breakthrough_Prize_in_Mathematics, And the citation of his work, https://breakthroughprize.org/?controller=Page&action=laureates&p=3&laureate_id=55.
67. E. Witten, Monopoles and four-manifolds, *Math. Res. Lett.* **1**, 769 (1994).
68. N. Seiberg and E. Witten, Electric-magnetic duality, monopole condensation, and confinement in $N = 2$ supersymmetric Yang–Mills theory, *Nucl. Phys. B* **426**, 19 (1994) [Erratum: *ibid.* **430**, 485 (1994)].
69. A. Jackson, A revolution in mathematics, <http://web.archive.org/web/20100426172959/http://www.ams.org/samplings/feature-column/mathnews-revolution>; D. Kotschick, Gauge theory is dead! — Long live gauge theory!, <http://www.ams.org/notices/199503/kotschick.pdf>; A. Scorpan, *The Wild World of 4-Manifolds* (American Mathematical Society, 2005).

70. E. Witten, An interpretation of classical Yang–Mills theory, *Phys. Lett. B* **77**, 394 (1978).
71. V. Volovich, Supersymmetric Yang–Mills theories and twistors, *Phys. Lett. B* **129**, 429 (1983), *Theor. Math. Phys.* **57**, 39 (1983); C. Devchand, An infinite number of continuity equations and hidden symmetries in supersymmetric gauge theories, *Nucl. Phys. B* **238**, 333 (1984); L. L. Chau, M-L. Ge and Z. Popowicz, Riemann–Hilbert transforms and Bianchi–Backlund transformations for the supersymmetric Yang–Mills fields, *Phys. Rev. Lett.* **52**, 1940 (1984); L. L. Chau, M-L. Ge and C. S. Lim, Constraints and equations of motion in supersymmetric Yang–Mills Theories, *Phys. Rev. D* **33**, 1056 (1986); Review by L-L Chau, Supersymmetric Yang–Mills fields as an integrable system and connections with other linear systems, in *Proceedings of the Workshop on Vertex Operators in Mathematical Physics*, Berkeley, 1983, eds. J. Lepowsky, S. Mandelstam and I. M. Singer (Springer, New York, 1984).
72. L. L. Chau and C. S. Lim, Geometrical constraints and equations of motion in extended supergravity, *Phys. Rev. Lett.* **56**, 294 (1986).
73. L. L. Chau, Linear systems and conservation laws of gravitational fields in four plus extended superspace, *Phys. Lett. B* **202**, 238 (1992).
74. N. Beisert *et al.*, Review of AdS/CFT integrability: An overview, *Lett. Math. Phys.* **99**, 3 (2012), and the references therein for all the review papers in the same volume.
75. L. D. Landau and E. M. Lifshitz, *Quantum Mechanics, Non-relativistic Theory*, Chapters XIV & XV, Vol. 3, *Theoretical Physics* (translated by J. B. Sykes and J. S. Bell, 1958).
76. G. F. Chew, *S-Matrix Theory of Strong Interactions*, Lecture Note & Reprint Series, (1961); *The Analytic S Matrix: A Basis for Nuclear Democracy*, (1966); G. F. Chew (author) and D. Pines (editor), *S-Matrix Theory Of Strong Interactions*, (2011).
77. L. Brink, Maximally supersymmetric Yang–Mills theory: The story of N=4 Yang–Mills theory, in this *Proceedings*.
78. C. M. Will (Guest ed.), Focus issue: Milestones of general relativity, *Class. Quantum Grav.* **32**, 120201 (2015), <http://iopscience.iop.org/0264-9381/page/Focus%20issue%20on%20Milestones%20of%20general%20relativity>.
79. W. T. Ni (ed.), One hundred years of general relativity: From genesis and empirical foundations to gravitational waves, cosmology and quantum gravity, Vol. 1 and 2 (World Scientific, Singapore, 2015), <http://www.worldscientific.com/worldscibooks/10.1142/9389>.
80. LIGO Scientific Collaboration and Virgo Collaboration, B. P. Abbott *et al.*, Observation of gravitational waves from a binary black hole merger, *Phys. Rev. Lett.* **116**, 061102 (Feb.12, 2016) and its pdf file at <https://physics.aps.org/featured-article-pdf/10.1103/PhysRevLett.116.061102>; and a report about it, <http://www.nytimes.com/2016/02/12/science/ligo-gravitational-waves-black-holes-einstein.html>.
81. In the opinion of many, including the author, it has been established that mathematics is the precise language of Nature. The author hopes that in some small measure this paper has helped to make the point that everyone, scientists and non-scientists, can participate in the excitement and the

enrichment found in the frontiers of scientific discovery, just as everyone can enjoy music and painting without being able to play a musical instrument and to paint. This point of view has guided the author in communicating physics (particle, condensed matter, and cosmology) to non-scientists since the late 1980s. For a sampling of the approach with good results, see the web page of the author’s 2005 Physics 10 for non-scientist undergraduates, <http://www.physics.ucdavis.edu/Classes/Physics10-B-W05/>.